arXiv:1507.01628v1 [cs.IT] 6 Jul 2015

NEW EXTREMAL BINARY SELF-DUAL CODES FROM A MODIFIED FOUR CIRCULANT CONSTRUCTION

ABIDIN KAYA, BAHATTIN YILDIZ, AND ABDULLAH PASA

Abstract. In this work, we propose a modified four circulant construction for self-dual codes and a bordered version of the construction using the properties of λ-circulant and λ-reverse circulant matrices. By using the constructions on $\mathbb{F}_2$, we obtain new binary codes of lengths 64 and 68. We also apply the constructions to the ring R_2 and considering the $\mathbb{F}_2$ and R_1-extensions, we obtain new singly-even extremal binary self-dual codes of lengths 66 and 68. More precisely, we find 3 new codes of length 64, 15 new codes of length 66 and 22 new codes of length 68. These codes all have weight enumerators with parameters that were not known to exist in the literature.

1. Introduction

Self-dual codes, especially over the binary field, have been a central topic of research in coding theory for a considerable time now. This interest is justified by the many combinatorial and algebraic objects that are related to self-dual codes. The recent studies on self-dual codes are intensified around classifying all extremal self-dual codes of given lengths. The possible weight enumerators of all extremal binary self-dual codes of lengths up to 100 are given in [4] and [6]. Many of these possible weight enumerators have parameters in them that fall in certain intervals. A lot of recent research on extremal self-dual codes has gone towards finding extremal self-dual codes with new parameters. [10], [11], [12], [13], [14], [15], [18] are examples of such works.

There are numerous constructions for extremal self-dual codes. The common theme is a computer search, usually using the magma computer algebra ([3]), over a reduced search field. There are different methods to reduce the search field. Some of the most common such methods use a special type of matrices called *circulant matrices*. An $n \times n$ circulant matrix is determined uniquely by its first row. So, searching through such matrices over an alphabet A, requires searching over $|A|^n$ matrices instead of $|A|^{n^2}$ matrices. Circulant matrices also have an important algebraic property in that they commute with respect to the matrix multiplication. These properties of circulant matrices have been used quite effectively in such popular constructions in literature as double circulant, bordered double circulant and four circulant constructions.

Another method of reducing the search field is to consider the above mentioned constructions over suitable rings that are endowed with orthogonality-preserving Gray maps. By considering the lifts of good binary self-dual codes over such rings, the search field can be reduced even further, giving way to many extremal codes with

---

2010 *Mathematics Subject Classification.* Primary 94B05; Secondary 94B99.

Key words and phrases. extremal self-dual codes, circulant matrices, reverse-circulant matrices, four circulant construction.

new parameters. This idea has been used quite effectively in the above-mentioned works.

In this work, we modify the four-circulant construction, using λ-circulant, reverse-circulant and λ-reverse-circulant matrices. Applying these modified constructions to three well-known alphabets, $\mathbb{F}_2$, $R_1 = \mathbb{F}_2 + u\mathbb{F}_2$ and $R_2 = \mathbb{F}_2 + u\mathbb{F}_2 + v\mathbb{F}_2 + uv\mathbb{F}_2$, we are able to find many extremal binary self-dual codes with new parameters, thus extending the database of all known such codes in the literature.

The rest of the work is organized as follows. In section 2, we give the preliminaries for our constructions. We recall the alphabets that we have mentioned above and we give a thorough background on circulant and reverse-circulant matrices. In section 3, we introduce a modified four-circulant construction, using λ-circulant and λ-reverse-circulant matrices as well as a bordered four circulant construction. In section 4, we apply the constructions to the alphabets $\mathbb{F}_2$, $R_1 = \mathbb{F}_2 + u\mathbb{F}_2$ and $R_2 = \mathbb{F}_2 + u\mathbb{F}_2 + v\mathbb{F}_2 + uv\mathbb{F}_2$, as a result of which we obtain many new extremal binary self-dual codes of lengths 64, 66 and 68. More explicitly, we construct 3 new codes of length 64, 15 new codes of length 66 and 22 new codes of length 68, adding these to the literature of known such codes. The results are tabulated for each length.

2. PRELIMINARIES

2.1. **Basics.** Let R be a finite ring. A linear *code* C of length n over R is an R-submodule of R^n. The elements of C are called *codewords.*

Let $\langle u, v\rangle$ be inner product of two codewords u and v in R^n which is defined as $\langle u, v\rangle = \sum_{i=1}^{n} u_i v_i$, where the operations are done in R^n. The *dual code* of a code C is $C^{\perp} = \{v \in R^n \mid \langle u, v\rangle = 0 \text{ for all } v \in C\}$. If $C \subseteq C^{\perp}$, C is called *self-orthogonal*, and C is *self-dual* if $C = C^{\perp}$.

The main case of interest for us is the case when $R = \mathbb{F}_2$, in which case we obtain the usual binary self-dual codes. Binary self-dual codes are called Type II if the weights of all codewords are multiples of 4 and Type I otherwise. Rains finalized the upper bound for the minimum distance d of a binary self-dual code of length n in [17] as $d \le 4\lfloor \frac{n}{24} \rfloor + 6$ if $n \equiv 22 \pmod{24}$ and $d \le 4\lfloor \frac{n}{24} \rfloor + 4$, otherwise. A self-dual binary code is called *extremal* if it meets the bound.

The ring family of R_k is introduced in [8] as follows:

$$R_k = \mathbb{F}_2[u_1, u_2, \ldots, u_k]/(u_i^2 = 0, u_i u_j = u_j u_i)$$

Leaving the details of these rings to the mentioned work, we just give basic properties of first two members of this family which are used in this work.

The ring R_1 that is $\mathbb{F}_2 + u\mathbb{F}_2$ is the first example of the ring family of R_k. This ring was introduced in [1] for constructing lattices and it has been studied quite extensively in the literature of coding theory. Second member of R_k is the ring R_2 that is $\mathbb{F}_2 + u\mathbb{F}_2 + v\mathbb{F}_2 + uv\mathbb{F}_2$. The ring R_2 was first introduced by Yildiz and Karadeniz in [20]. Unlike the ring R_2, the ring R_1 is finite chain ring. Moreover, the ring R_k is not finite chain ring for $k \ge 2$.

A linear distance preserving map from R_1^n to $\mathbb{F}_2^{2n}$ is described in [5] in terms of vectors as:

$$\phi_1(\bar{a} + u\bar{b}) = (\bar{b}, \bar{a} + \bar{b}),$$

where $\bar{a}, \bar{b} \in \mathbb{F}_2^{2n}$.

In [20], the Lee weight was defined as the Hamming weight of the image under a Gray map which was extended from that in [5]. Specifically, the following map for R_2 is given in [20] as follows:

$$\phi_2(\bar{a}+u\bar{b}+v\bar{c}+uv\bar{d}) = (\bar{d}, \bar{c}+\bar{d}, \bar{b}+\bar{d}, \bar{a}+\bar{b}+\bar{c}+\bar{d}),$$

where $\bar{a}, \bar{b}, \bar{c}, \bar{d} \in \mathbb{F}_2^n$.

One of the important properties of the Gray map that was given for this family of rings was its orthogonality-preserving property. This gives rise to the following important lemma which was given as Theorem 4 in [9]:

Lemma 2.1. *Let C be a self-dual code over R_k of length n then $\phi_k(C)$ is a binary self-dual code of length $2^k n$. If C is Type II code then $\phi_k(C)$ is Type II and if C is Type I then $\phi_k(C)$ is Type I.*

2.2. **The Circulant and Reverse-Circulant Matrices.**

Since our constructions rely heavily on circulant and reverse-circulant matrices, we give some of their properties in this subsection. Most of what is explained here can be found in abstract algebra sources. However, we wanted to put together some of these in a complete form.

With R a commutative ring with identity, let σ be the permutation on R^n that corresponds to the right shift, i.e.

$$\sigma(a_1, a_2, \ldots, a_n) = (a_n, a_1, \ldots, a_{n-1}). \tag{2.1}$$

A circulant matrix is a square matrix where each row is a right-circular shift of the previous row. In other words, if $\overline{r}$ is the first row, a typical circulant matrix is of the form

$$\left[\begin{array}{c} \overline{r} \\ \hline \sigma(\overline{r}) \\ \hline \sigma^2(\overline{r}) \\ \hline \vdots \\ \hline \sigma^{n-1}(\overline{r}) \end{array}\right]. \tag{2.2}$$

It is clear that, with T denoting the permutation matrix corresponding to the n-cycle $(123...n)$, a circulant matrix with first row $(a_1, a_2, \ldots, a_n)$ can be expressed as a polynomial in T as:

$$a_1 I_n + a_2 T + a_3 T^2 + \cdots + a_n T^{n-1}.$$

Since T satisfies $T^n = I_n$, this shows that circulant matrices commute.

A reverse-circulant matrix is a square matrix where each row is a left-circular shift of the previous row. It is clear to see that if $\overline{r}$ is the first row, a reverse-circulant matrix is of the form

$$\left[\begin{array}{c} \overline{r} \\ \hline \sigma^{-1}(\overline{r}) \\ \hline \sigma^{-2}(\overline{r}) \\ \hline \vdots \\ \hline \sigma^{-(n-1)}(\overline{r}) \end{array}\right]. \tag{2.3}$$

The following lemma recaptures some of the basic properties of circulant and reverse-circulant matrices that can easily be verified:

Lemma 2.2. ***(i)*** *Circulant matrices commute under matrix multiplication. Reverse-circulant matrices do not necessarily commute under multiplication.*

(ii) *Reverse-circulant matrices are symmetric.*

(iii) *Suppose that A is a circulant matrix and D is the backdiagonal matrix with 1 on its backdiagonal and 0 elsewhere. Then both AD and DA are reverse circulant.*

Since $D^2 = I_n$, D is its own inverse. This means that from (iii) of the previous lemma we get the following corollary:

Corollary 2.3. *Let A be a reverse-circulant matrix. Then there exists unique circulant matrices B and C such that $A = BD = DC$.*

We prove the following important lemma using the properties above:

Lemma 2.4. ***(i)*** *Circulant matrices are closed under matrix multiplication. The product of two reverse-circulant matrices is circulant.*

(ii) *If A is a circulant matrix and B is a reverse-circulant matrix, then AB and BA are both reverse-circulant matrices that are not necessarily the same matrices.*

Proof. **(i)** If A_1 and A_2 are circulant matrices then they correspond to two polynomials in T with $T^n = I_n$. It is clear that their product is also a polynomial in T of degree at most $n-1$.

Now suppose that B_1 and B_2 are two reverse-circulant matrices. By Corollary 2.3, we see that there exists circulant matrices A_1 and A_2 such that $B_1 = A_1 D$ and $B_2 = DA_2$. Since $D^2 = I_n$, we get

$$B_1 B_2 = (A_1 D)(DA_2) = A_1(D^2)A_2 = A_1 A_2,$$

which is circulant by the first part.

(ii) Let C_1 and C_2 be circulant matrices such that $B = DC_1 = C_2 D$ by Corollary 2.3. Then $AB = A(C_2 D) = (AC_2)D$, which is reverse-circulant by Lemma 2.2 (iii) and the first part. Similarly, $BA = (DC_1)A = D(C_1 A)$ is reverse-circulant. □

Let λ be a unit in R. The circulant and reverse-circulant matrices can be extended to the so-called λ-circulant and λ-reverse-circulant matrices. By σ_λ, we mean the map that acts on R^n as

$$\sigma_\lambda(a_1, a_2, \ldots, a_n) = (\lambda a_n, a_1, \ldots, a_{n-1}). \tag{2.4}$$

A λ-circulant matrix is then a square matrix where each row is obtained from the previous row by applying the map σ_λ. In other words, a typical λ-circulant matrix is of the form

$$\left[\begin{array}{c} \overline{r} \\ \hline \sigma_\lambda(\overline{r}) \\ \hline \sigma_\lambda^2(\overline{r}) \\ \hline \vdots \\ \hline \sigma_\lambda^{n-1}(\overline{r}) \end{array}\right]. \tag{2.5}$$

Let T_λ be the matrix that is obtained from the permutation matrix T by multiplying the $(n,1)$-entry by λ. Then a λ-circulant matrix with the first row $(a_1, a_2, \ldots, a_n)$ can be expressed as

$$a_1 I_n + a_2 T_\lambda + a_3 T_\lambda^2 + \cdots + a_n T_\lambda^{n-1}$$

with $T_\lambda^n = \lambda I_n$. This shows that λ-circulant matrices are closed under multiplication and that they commute with each other as well.

By ρ_λ, let us define the analogous left λ-shift on R^n, namely:

$$\rho_\lambda(a_1, a_2, \ldots, a_n) = (a_2, a_3, \ldots, a_n, \lambda a_1). \tag{2.6}$$

From the definitions, we can easily observe the following identities between σ_λ and ρ_λ

$$\sigma_{\lambda^{-1}} \circ \rho_\lambda = \rho_\lambda \circ \sigma_{\lambda^{-1}} = 1. \tag{2.7}$$

A λ-reverse-circulant matrix is defined as a square matrix where each row is obtained from the previous row by applying the map ρ_λ. In other words, a typical λ-reverse-circulant matrix is of the form

$$\begin{bmatrix} \overline{r} \\ \hline \rho_\lambda(\overline{r}) \\ \hline \rho_\lambda^2(\overline{r}) \\ \hline \vdots \\ \hline \rho_\lambda^{n-1}(\overline{r}) \end{bmatrix}. \tag{2.8}$$

Similar to the properties of circulant and reverse-circulant matrices we have the following results for λ-circulant and λ-reverse-circulant matrices.

Lemma 2.5. ***(i)*** *The product of two λ-circulant matrices is again λ-circulant while the product of a λ-circulant and γ-circulant matrix is not necessarily circulant if $\gamma \neq \lambda$.*

(ii) *λ-circulant matrices commute under multiplication for the same λ.*

(iii) *λ-reverse-circulant matrices are symmetric.*

(iv) *With D denoting the backdiagonal matrix with 1's on its backdiagonal, and A denoting a λ-circulant matrix, we have: AD is a λ-reverse-circulant matrix while DA is a λ^{-1}-reverse-circulant matrix.*

(v) *For any λ-reverse-circulant matrix A, there exists a unique λ-circulant matrix B and a unique λ^{-1}-circulant matrix C such that $A = BD = DC$.*

We prove the following result, which is analogous to Lemma 2.4:

Theorem 2.6. *Suppose A is a λ-circulant matrix and B be a λ-reverse-circulant matrix. Then AB is a λ-reverse-circulant matrix for all units $\lambda \in R$. If $\lambda^2 = 1$, then BA is also λ-reverse-circulant.*

Proof. By Lemma 2.5-(v), let C_1 be the λ-circulant matrix such that $B = C_1D$. Then $AB = A(C_1D) = (AC_1)D$. Since both A and C_1 are λ-circulant, AC_1 is λ-circulant, and so by Lemma 2.5-(iv), $(AC_1)D$ is λ-reverse-circulant.

For the second part, assume that $\lambda^2 = 1$, $\lambda^{-1} = \lambda$. Again by the previous lemma, let C_2 be the $\lambda^{-1} = \lambda$-circulant matrix such that $B = DC_2$. Then $BA = (DC_2)A = D(C_2A)$. Similar to the previous case, C_2A is λ-circulant, and so by the same lemma $BA = D(C_2A)$ is $\lambda^{-1} = \lambda$-reverse-circulant. □

3. Constructions

The four circulant construction were introduced in [2] as;

Let A and B be $n \times n$ circulant matrices over $\mathbb{F}_p$ such that $AA^T + BB^T = -I_n$ then the matrix

$$G = \left[I_{2n} \,\middle|\, \begin{matrix} A & B \\ -B^T & A^T \end{matrix} \right]$$

generates a self-dual code over $\mathbb{F}_p$.

Recently, the construction is applied on $\mathbb{F}_2 + u\mathbb{F}_2$ in [11], which yielded to new binary self-dual codes.

In the following we give a modified version of the construction, which works for any commutative Frobenius ring;

Theorem 3.1. *Let λ be a unit of the commutative Frobenius ring R, A be a λ-circulant matrix and B be a λ-reverse-circulant matrix with $AA^T + BB^T = -I_n$ then the matrix*

$$G = \left[I_{2n} \;\middle|\; \begin{array}{cc} A & B \\ -B & A \end{array} \right]$$

generates a self-dual code $\mathcal{C}$ over R.

Proof. Let $X = \begin{bmatrix} A & B \\ -B & A \end{bmatrix}$, it is enough to show that $XX^T = -I_{2n}$. We have

$$XX^T = \begin{bmatrix} -I_n & -AB^T + BA^T \\ -BA^T + AB^T & -I_n \end{bmatrix}.$$

So, we need to verify that $-AB^T + BA^T = 0$. By Lemma 2.5-(iii) B is a symmetric matrix so $AB^T = AB$ and by Theorem 2.6 AB is a λ-reverse-circulant matrix and it is also symmetric by Lemma 2.5-(iii). It follows that $AB^T = BA^T$, which implies $GG^T = 0$. Since the ring R is Frobenius, the code $\mathcal{C}$ is self-dual. $\square$

By applying Theorem 3.1 on $\mathbb{F}_2$ we obtain the codes listed in Table 1. Some examples of codes over R_2 are given in Table 2.

For the rest of the manuscript let $\boldsymbol{J}_n$ denote the $n \times n$ matrix with all entries equal to 1.

Lemma 3.2. *Let A be a λ-circulant matrix over R_k, B be a reverse-circulant matrix and $r_A = (a_1, a_2, \ldots, a_n)$ and $r_B = (b_1, b_2, \ldots, b_n)$ denote the first rows of A and B, respectively. Let S_{r_A} and S_{r_B} denote the sum of the components of r_A and r_B, respectively, i.e.$S_{r_A} = \sum_{i=1}^{n} a_i$. Then, $AA^T + BB^T = I_n + \boldsymbol{J}_n$ implies S_{r_A} and S_{r_B} are both units or both non-units in R_k.*

Proof. $AA^T + BB^T = I_n + \boldsymbol{J}_n$ implies $\langle r_A, r_A\rangle + \langle r_B, r_B\rangle = 0$, i.e. $\sum_{i=1}^{n} a_i^2 + \sum_{i=1}^{n} b_i^2 = 0$. Then, $\sum_{i=1}^{n} a_i^2 = (S_{r_A})^2$ and $\sum_{i=1}^{n} b_i^2 = (S_{r_B})^2$ since $char\,(R_k) = 2$. Hence, $(S_{r_A})^2 = (S_{r_B})^2$ and the result follows from the fact $(\text{unit})^2 = 1$, $(\text{non-unit})^2 = 0$ for elements of R_k. $\square$

The following Theorem is a bordered version of the construction in Theorem 3.1, which generates self-dual codes over the family of rings R_k.

Theorem 3.3. *Let n be an odd number, A be a circulant matrix or order n and B be a reverse-circulant matrix of order n with $S_{r_A} = S_{r_B}$, which is a unit and $AA^T + BB^T = I_n + \boldsymbol{J}_n$. Let x be a unit and y be a non-unit in R_k. Let $z = xS_{r_A}$ and $t = yS_{r_B}$ then the matrix*

$$G = \left[I_{2n+2} \;\middle|\; \begin{array}{cccc} 1 & 1 & \boldsymbol{x} & \boldsymbol{y} \\ 1 & 1 & \boldsymbol{y} & \boldsymbol{x} \\ \boldsymbol{z}^T & \boldsymbol{t}^T & A & B \\ \boldsymbol{t}^T & \boldsymbol{z}^T & B & A \end{array} \right]$$

generates a self-dual code $\mathcal{C}$ over R_k where $\boldsymbol{x} = (x, x, \ldots, x)$ denotes the all-x vector, similarly for $\boldsymbol{y}$, $\boldsymbol{z}$ and $\boldsymbol{t}$.

Proof. Let g_i denote the i-th row of G. $\langle g_1, g_1\rangle = 1+n\left(x^2+y^2\right) = 0$ since n is odd, $x^2 = 1$ and $y^2 = 0$. It is obvious that $\langle g_1, g_2\rangle = 0$. By the choice of z and t we have $\langle g_1, g_i\rangle = 0$ for $3 \leq i \leq 2n+2$. Similarly g_2 is orthogonal to itself and the other rows. Since z is a unit and t is a non-unit we have $z^2 = 1$ and $t^2 = 0$ which implies $\langle g_i, g_i\rangle = 1+z^2+t^2 = 0$ for $3 \leq i \leq 2n+2$. $AA^T+BB^T = I_n+\boldsymbol{J}_n$ and $z^2+t^2 = 1$ implies $\langle g_i, g_j\rangle = 0$ whenever $3 \leq i < j \leq n+2$ or $n+3 \leq i < j \leq 2n+2$. By Lemma 2.4-(ii) AB is a reverse-circulant matrix and by Lemma 2.2-(ii) reverse-circulant matrices are symmetric which implies $B = B^T$ and therefore $AB^T+BA^T = 0$. Together with $zt+tz = 0$ this implies $\langle g_i, g_j\rangle = 0$ whenever $3 \leq i \leq n+2$ or $n+3 \leq j \leq 2n+2$. Hence, $\mathcal{C}$ is a self-dual code over R_k. $\square$

Example 3.4. Let $\mathcal{C}$ be the code over R_1 generated by

$$\left[\begin{array}{c|cccc} & 1 & 1 & \mathbf{1} & \boldsymbol{u} \\ & 1 & 1 & \boldsymbol{u} & \mathbf{1} \\ I_{16} & (\mathbf{1}+\boldsymbol{u})^T & \boldsymbol{u}^T & A & B \\ & \boldsymbol{u}^T & (\mathbf{1}+\boldsymbol{u})^T & B & A \end{array}\right]$$

where $r_A = (u,0,1,1,u,1,u)$ and $r_B = (0,0,0,1,u,u,u)$ then $S_{r_A} = S_{r_B} = 1+u$ and $AA^T+BB^T = I_7+\boldsymbol{J}_7$. Thus, $\mathcal{C}$ is a self-dual code by Theorem 3.3. The binary image $\phi_1(\mathcal{C})$ is a self-dual Type II $[64,32,12]_2$-code.

The codes listed in Table 3 are obtained by applying Theorem 3.3 on $\mathbb{F}_2$.

4. New binary self-dual codes of lengths 64, 66 and 68

In this section, we apply the constructions given in Section 3. By using Theorem 3.1 on $\mathbb{F}_2$ we were able to construct 4 new codes of length 68. Three new extremal binary self-dual codes of length 64 are constructed as an application of Theorem 3.3 on $\mathbb{F}_2$. Moreover, by considering the extensions of the codes new codes of length 66 and 68 are constructed. The following theorem is used for extensions:

Theorem 4.1. ([7]) *Let $\mathcal{C}$ be a self-dual code over R of length n and $G = (r_i)$ be a $k \times n$ generator matrix for $\mathcal{C}$, where r_i is the i-th row of G, $1 \leq i \leq k$. Let c be a unit in R such that $c^2 = 1$ and X be a vector in R^n with $\langle X, X\rangle = 1$. Let $y_i = \langle r_i, X\rangle$ for $1 \leq i \leq k$. Then the following matrix*

$$\left(\begin{array}{cc|c} 1 & 0 & X \\ \hline y_1 & cy_1 & r_1 \\ \vdots & \vdots & \vdots \\ y_k & cy_k & r_k \end{array}\right),$$

generates a self-dual code $\mathcal{C}'$ over R of length $n+2$.

4.1. New extremal binary self-dual codes of length 64.

There are two possibilities for the weight enumerators of extremal singly-even $[64,32,12]_2$ codes ([4]):

$$\begin{aligned} W_{64,1} &= 1+(1312+16\beta)y^{12}+(22016-64\beta)y^{14}+\cdots, \ 14 \leq \beta \leq 284, \\ W_{64,2} &= 1+(1312+16\beta)y^{12}+(23040-64\beta)y^{14}+\cdots, \ 0 \leq \beta \leq 277. \end{aligned}$$

Together with the ones constructed in [10, 11, 18], codes exist with weight enumerators β =14, 18, 22, 25, 32, 36, 39, 44, 46, 53, 60, 64 in $W_{64,1}$ and for β =0, 1, 2, 4, 5, 6, 8, 9, 10, 12, 13, 14, 16, 17, 18, 20, 21, 22, 23, 24, 25, 28, 29, 30, 32, 33, 36,

Table 1. Extremal binary self-dual Type I codes of length 64 by Theorem 3.1 on $\mathbb{F}_2$.

$\mathcal{B}_{64,i}$	r_A	r_B	β in $W_{64,2}$	$\lvert Aut\left(\mathcal{B}_{64,i}\right)\rvert$
$\mathcal{B}_{64,1}$	(0101110001100111)	(1011010101010100)	0	2^5
$\mathcal{B}_{64,2}$	(1010101011110101)	(0110110001001001)	8	2^5
$\mathcal{B}_{64,3}$	(0111000010111110)	(0011001111010010)	16	2^5
$\mathcal{B}_{64,4}$	(1011001001100101)	(0110100101000000)	24	2^5
$\mathcal{B}_{64,5}$	(1100110010100010)	(1110010010111110)	32	2^5
$\mathcal{B}_{64,6}$	(1000111110101000)	(0110100011011101)	40	2^5
$\mathcal{B}_{64,7}$	(0100010000111100)	(1011101010000001)	48	2^5
$\mathcal{B}_{64,8}$	(1111011010000100)	(1101010101100011)	56	2^5
$\mathcal{B}_{64,9}$	(1000110110110001)	(1011001001101011)	64	2^6
$\mathcal{B}_{64,10}$	(0101110111001111)	(0001000111000110)	72	2^5

37, 38, 40, 41, 44, 48, 51, 52, 56, 58, 64, 72, 80, 88, 96, 104, 108, 112, 114, 118, 120 and 184 in $W_{64,2}$.

In this work, we construct the codes with weight enumerators β =29, 59, 74 in $W_{64,1}$.

By Theorem 3.1 we obtain extremal binary self-dual codes of length 64 that are given in Table 1.

In order to fit the upcoming tables we use hexadecimal number sytem. The one-to-one correspondence between hexadecimals and binary 4 tuples is as follows:

$$\begin{aligned}
0 &\leftrightarrow 0000,\ 1 \leftrightarrow 0001,\ 2 \leftrightarrow 0010,\ 3 \leftrightarrow 0011,\\
4 &\leftrightarrow 0100,\ 5 \leftrightarrow 0101,\ 6 \leftrightarrow 0110,\ 7 \leftrightarrow 0111,\\
8 &\leftrightarrow 1000,\ 9 \leftrightarrow 1001,\ A \leftrightarrow 1010,\ B \leftrightarrow 1011,\\
C &\leftrightarrow 1100,\ D \leftrightarrow 1101, E \leftrightarrow 1110,\ F \leftrightarrow 1111.
\end{aligned}$$

To express the elements of R_2 we use the ordered basis $\{uv, v, u, 1\}$. As an application of Theorem 3.1 on R_2 binary self-dual Type I codes of length 64 are constructed, which are listed in Table 2.

Remark 4.2. Note that the first codes with weight enumerators β =1, 5, 13, 17, 21, 25, 29, 33 and 80 in $W_{64,2}$ are recently constructed in [10, 11]. Those are reconstructed in Table 2.

The construction in Theorem 3.3 yields extremal binary self-dual codes of length 64. Those are listed in Table 3, codes with weight enumerators β =29, 59 and 74 in $W_{64,1}$ are constructed for the first time.

4.2. New extremal binary self-dual codes of length 66.

A self-dual $[66,33,12]_2$-code has a weight enumerator in one of the following forms ([6]):

$$\begin{aligned}
W_{66,1} &= 1+(858+8\beta)\,y^{12}+(18678-24\beta)\,y^{14}+\cdots,\ \text{where } 0\le\beta\le 778,\\
W_{66,2} &= 1+1690y^{12}+7990y^{14}+\cdots\\
\text{and } W_{66,3} &= 1+(858+8\beta)\,y^{12}+(18166-24\beta)\,y^{14}+\cdots,\ \text{where } 14\le\beta\le 756,
\end{aligned}$$

For a list of known codes in $W_{66,1}$ we refer to [11, 15, 19] and codes with weight enumerator $W_{66,2}$ exist. Together with the ones constructed in [12] the existence of the codes is known for β =28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 43, 44, 45,

TABLE 2. Extremal self-dual $[64, 32, 12]_2$ codes as Gray images by Theorem 3.1 on R_2.

$\mathcal{D}_{64,i}$	λ	r_A	r_B	β in $W_{64,2}$	$\lvert Aut(\mathcal{D}_{64,i})\rvert$
$\mathcal{D}_{64,1}$	5	$(6, 5, A, E)$	$(D, 7, D, 5)$	0	2^5
$\mathcal{D}_{64,2}$	B	$(1, 6, 3, B)$	$(1, 8, 7, 4)$	1	2^3
$\mathcal{D}_{64,3}$	7	$(9, 7, 9, D)$	$(5, C, A, A)$	4	2^3
$\mathcal{D}_{64,4}$	7	$(5, 2, 3, 9)$	$(7, 4, 3, 8)$	5	2^3
$\mathcal{D}_{64,5}$	7	$(6, F, A, B)$	$(C, 9, 3, 1)$	8	2^3
$\mathcal{D}_{64,6}$	D	$(9, E, D, B)$	$(F, A, 5, 0)$	9	2^3
$\mathcal{D}_{64,7}$	7	$(4, 9, F, 9)$	$(7, 8, B, 6)$	12	2^4
$\mathcal{D}_{64,8}$	D	$(F, E, 3, 3)$	$(D, 2, B, 4)$	13	2^3
$\mathcal{D}_{64,9}$	3	$(3, 3, 5, B)$	$(8, 4, 2, 9)$	16	2^5
$\mathcal{D}_{64,10}$	B	$(D, 4, 7, 9)$	$(D, 6, B, 0)$	17	2^3
$\mathcal{D}_{64,11}$	D	$(3, 7, 9, B)$	$(3, 2, E, E)$	20	2^3
$\mathcal{D}_{64,12}$	7	$(E, 7, 0, 4)$	$(B, 3, 5, 7)$	21	2^3
$\mathcal{D}_{64,13}$	7	$(C, 1, 1, 9)$	$(5, 2, F, 8)$	24	2^4
$\mathcal{D}_{64,14}$	7	$(8, B, 6, 2)$	$(1, 1, D, F)$	25	2^3
$\mathcal{D}_{64,15}$	B	$(C, D, 0, 3)$	$(D, E, 7, 5)$	28	2^5
$\mathcal{D}_{64,16}$	B	$(6, 7, C, 4)$	$(F, D, 9, D)$	29	2^3
$\mathcal{D}_{64,17}$	3	$(5, 4, 1, 4)$	$(7, B, 7, 6)$	32	2^5
$\mathcal{D}_{64,18}$	D	$(8, 5, 4, 2)$	$(1, B, 5, 1)$	33	2^3
$\mathcal{D}_{64,19}$	B	$(9, 9, C, 3)$	$(8, 1, A, F)$	36	2^3
$\mathcal{D}_{64,20}$	5	$(E, A, D, 6)$	$(F, 3, B, D)$	48	2^5
$\mathcal{D}_{64,21}$	D	$(6, 9, 0, 3)$	$(A, 9, 3, 1)$	64	2^5
$\mathcal{D}_{64,22}$	5	$(A, 9, D, 1)$	$(F, 8, 5, E)$	80	2^7

TABLE 3. Extremal self-dual Type I $[64, 32, 12]_2$ codes by Theorem 3.3 on $\mathbb{F}_2$.

$\mathcal{C}_{64,i}$	r_A	r_B	β in $W_{64,1}$	$\lvert Aut(\mathcal{C}_{64,i})\rvert$
$\mathcal{C}_{64,1}$	(001101000000011)	(011000010011011)	14	$2^2 \times 3 \times 5$
$\mathcal{C}_{64,2}$	(010001101111110)	(111111100011110)	14	$2 \times 3 \times 5$
$\mathcal{C}_{64,3}$	(001101111000010)	(110010110110011)	**29**	$2 \times 3 \times 5$
$\mathcal{C}_{64,4}$	(111010001101101)	(100101000001111)	44	$2 \times 3 \times 5$
$\mathcal{C}_{64,5}$	(101000110101111)	(000000000011100)	44	$2^2 \times 3 \times 5$
$\mathcal{C}_{64,6}$	(101101011101111)	(001000001110001)	**59**	$2 \times 3 \times 5$
$\mathcal{C}_{64,7}$	(011000100111111)	(011000000000010)	**74**	$2^2 3 \times 5$

46, 47, 48, 49, 50, 51, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 66, 67, 70, 71, 73, 74, 75, 76, 77, 78, 79 and 80 in $W_{66,3}$.

In this work, we construct 15 codes with new weight enumerators in $W_{66,3}$. More precisely, the codes with weight enumerators β =46, 52, 53, 61, 64, 81, 82, 83, 84, 85, 86, 87, 88, 90 and 92 in $W_{66,3}$ are constructed for the first time in the literature. The codes that are obtained by applying Theorem 4.1 are listed in Table 4 and Table 5.

Table 4. New codes in $W_{66,3}$ by Theorem 4.1 $\mathbb{F}_2$. (10 codes)

$\mathcal{C}_i$	X	β in $W_{66,3}$
$\mathcal{C}_{64,5}$	$\left(11001101110010001101111011101100\mathbf{1}^{\mathbf{32}}\right)$	**52**
$\mathcal{C}_{64,4}$	$\left(11110101011010111010110100000001\mathbf{1}^{\mathbf{32}}\right)$	**61**
$\mathcal{C}_{64,5}$	$\left(00100101011000001000111010101100\mathbf{1}^{\mathbf{32}}\right)$	**64**
$\mathcal{C}_{64,7}$	$\left(11110100100100011110100101100101\mathbf{1}^{\mathbf{32}}\right)$	**81**
$\mathcal{C}_{64,7}$	$\left(11110011010001000000011101011110\mathbf{1}^{\mathbf{32}}\right)$	**83**
$\mathcal{C}_{64,7}$	$\left(00101011111010100110001001011110\mathbf{1}^{\mathbf{32}}\right)$	**84**
$\mathcal{C}_{64,7}$	$\left(00101111000111010000101010111101\mathbf{1}^{\mathbf{32}}\right)$	**85**
$\mathcal{C}_{64,7}$	$\left(11001000011100000101010011000110\mathbf{1}^{\mathbf{32}}\right)$	**87**
$\mathcal{C}_{64,7}$	$\left(11111000010000100011011010100101\mathbf{1}^{\mathbf{32}}\right)$	**90**
$\mathcal{C}_{64,7}$	$\left(01010110000100110011000110000011\mathbf{1}^{\mathbf{32}}\right)$	**92**

Table 5. New codes in $W_{66,3}$ by Theorem 4.1 $\mathbb{F}_2$. (5 codes)

$\mathcal{C}_{64,i}$	X	β
$\mathcal{C}_{64,3}$	(1100110010100000010111010111000010110000110011010000111001101100)	**46**
$\mathcal{C}_{64,5}$	(0010001010110101110110100110011000110110101100100000110000111101)	**53**
$\mathcal{C}_{64,7}$	(0000101100110000000001100100101110100001010010101111110011001001)	**82**
$\mathcal{C}_{64,7}$	(0011000101011100001001101011011000101100110100101110000011100010)	**86**
$\mathcal{C}_{64,7}$	(1110110000111111101101111011001110101101010101100100001101111001)	**88**

4.3. **New extremal binary self-dual codes of length 68.** The weight enumerator of a self-dual $[68,34,12]_2$ code is in one of the following forms ([6]):

$$
\begin{aligned}
W_{68,1} &= 1+(442+4\beta)\,y^{12}+(10864-8\beta)\,y^{14}+\cdots,\\
W_{68,2} &= 1+(442+4\beta)\,y^{12}+(14960-8\beta-256\gamma)\,y^{14}+\cdots
\end{aligned}
$$

where β and γ are parameters. Recently, 32 new codes are obtained in [12] and 178 new codes including the first examples with $\gamma=3$ in $W_{68,2}$ are obtained in [13]. For a list of known codes in $W_{68,1}$ we refer to [19, 15]. Recently, new codes in $W_{68,2}$ are obtained in [13, 12, 15, 19] together with these, codes exist for $W_{68,2}$ when

$$
\begin{aligned}
\gamma &= 0,\ \beta=11,22,33,44,\ldots,154,165,187,209,231,303\\
\text{or } \beta &\in \{2m \mid m=17,\ 20,\ 88,\ 99,\ 102,\ 110,\ 119,\ 136,\ 165 \text{ or } 78\le m\le 86\}\,;\\
\gamma &= 1,\ \beta=49,57,59,\ldots,160 \text{ or } \beta\in\{2m \mid m=27,\ 28,\ 29,\ 95,\ 96 \text{ or } 81\le m\le 89\}\,;\\
\gamma &= 2,\ \beta=65,69,71,77,81,159,186 \text{ or } \beta\in\{2m \mid 30\le m\le 68,\ 70\le m\le 91\} \text{ or}\\
\beta &\in \{2m+1 \mid 42\le m\le 69,\ 71\le m\le 77\}\,;\\
\gamma &= 3,\ \beta=101,107,117,123,127,133,137,141,145,147,149,153,159,193 \text{ or}\\
\beta &\in \{2m \mid m=44{,}45{,}47{,}48{,}50{,}51{,}52{,}54{,}\ldots{,}61{,}63{,}\ldots{,}66{,}68{,}\ldots{,}72{,}74{,}77{,}\ldots{,}84{,}86{,}\ldots{,}90{,}94{,}98\}\,;\\
\gamma &= 4,\ \beta\in\{2m \mid m=43{,}48{,}49{,}51{,}52{,}54{,}55{,}56{,}58{,}60{,}61{,}62{,}64{,}65{,}67{,}\ldots{,}71{,}75{,}\ldots{,}78{,}80{,}87{,}97\}\,;\\
\gamma &= 6 \text{ with } \beta\in\{2m \mid m=69,\ 77,\ 78,\ 79,\ 81,\ 88\}\,.
\end{aligned}
$$

We construct the codes with weight enumerators in $W_{68,2}$ for $\gamma=0$ and $\beta=$17, 155, 157, 187, 221 and 255; $\gamma=3$ and $\beta=$103, 105, 115, 119, 121, 124, 125, 129, 131, 134, 150, 178, 182, 184, 190 and 194.

TABLE 6. New codes in $W_{68,2}$ by Theorem 3.1 on $\mathbb{F}_2$. (4 codes)

$\mathcal{C}_{68,i}$	r_A	r_B	β in $W_{68,2}$	$\lvert Aut\left(\mathcal{C}_{68,i}\right)\rvert$
$\mathcal{C}_{68,1}$	(01111110101111011)	(11001000101001011)	**17**	2×17
$\mathcal{C}_{68,2}$	(11110001011001010)	(11010100001011010)	**187**	2×17
$\mathcal{C}_{68,3}$	(00110001101111011)	(01000010000000100)	**221**	2×17
$\mathcal{C}_{68,4}$	(11010010110010011)	(10100001001111100)	**255**	2×17

TABLE 7. New extremal binary self-dual codes of length 68 with $\gamma = 3$ in $W_{68,2}$ by Theorem 4.1 on R_1(16 codes)

Code	c	X	β in $W_{68,2}$
$\mathcal{D}_{64,10}$	1	$(1u1001030u3103111u3130u01u0u0331)$	**103**
$\mathcal{D}_{64,10}$	3	$(303u0101uu3301113u31300u1000u113)$	**105**
$\mathcal{D}_{64,10}$	1	$(3u1u03u3u03303331u313u0u30uuu331)$	**115**
$\mathcal{D}_{64,10}$	1	$(3010u1u10u1103313u111u0u3uu00113)$	**119**
$\mathcal{D}_{64,10}$	3	$(301001030u1303131u31100u3u000133)$	**121**
$\mathcal{D}_{64,10}$	3	$(uuu101u303u3u11uu3u1003uu1u1001u)$	**124**
$\mathcal{D}_{64,10}$	1	$(1u1uu3030u11u3133u3330uu3u00u333)$	**125**
$\mathcal{D}_{64,10}$	3	$(1u1003u10u33u313303330u01000u111)$	**129**
$\mathcal{D}_{64,10}$	3	$(1u10u1030u13u31330331u001uu0u111)$	**131**
$\mathcal{D}_{64,10}$	3	$(1u303011uu01000u10303133u0u10u33)$	**134**
$\mathcal{D}_{64,10}$	1	$(000101u10103u110u1030u30u101uu30)$	**150**
$\mathcal{D}_{64,22}$	1	$(3u000103031u00u30uu03u30u0uu11u1)$	**178**
$\mathcal{D}_{64,22}$	1	$(3u0uu3u3u13uuuu3uuuu1u3uuu0031u3)$	**182**
$\mathcal{D}_{64,22}$	3	$(u1100u001uu0uuu311331u101u03111u)$	**184**
$\mathcal{D}_{64,22}$	1	$(1uu00301u33uuu03u00u3u1u0u0031u3)$	**190**
$\mathcal{D}_{64,22}$	1	$(30000101013u00u100003u100u001303)$	**194**

By using Theorem 3.1 we were able to obtain codes with weight enumerators $\gamma = 0$ and $\beta =$**17**, 34, 51, 68, 85, 102, 119, 136, 153, 170, **187**, 204, **221**, 238, **255**, 272 for $W_{68,2}$. In order to save space we only list the new codes in Table 6.

Example 4.3. Let $\mathcal{C}_1$ and $\mathcal{C}_2$ be the R_1-extensions of $\varphi_u\left(\mathcal{D}_{64,21}\right)$ with respect to Theorem respectively with $c_1 = 1 + u, c_2 = 1$ and

$$\begin{aligned}
X_1 &= (3,u,0,0,0,0,1,u,3,0,3,u,1,1,0,0,u,1,1,0,1,3,1,u,1,3,0,u,0,0,3,3) \\
X_2 &= (1,u,u,0,0,u,1,0,3,0,3,0,1,3,u,0,u,1,1,u,3,3,1,u,1,1,u,0,u,u,1,1).
\end{aligned}$$

Then, $\varphi\left(\mathcal{C}_1\right)$ and $\varphi\left(\mathcal{C}_2\right)$ are extremal binary self-dual codes of length 68 with weight enumerators respectively $\gamma = 0, \beta = 155$ and $\gamma = 0, \beta = 157$ in $W_{68,2}$.

The codes over R_2 in Table 2 are mapped to R_1 and by applying the extension theorem we were able to obtain 16 new codes of length 68 as binary images of the extensions. The codes are listed in Table 7.

Theorem 4.4. *The existence of extremal self-dual binary codes is known for* 15 *parameters in* $W_{64,1}$; 57 *parameters in* $W_{66,3}$ *and* 465 *parameters in* $W_{68,2}$.

Remark 4.5. The binary generator matrices of the constructed new codes are available online at [16].

References

- [1] C. Bachoc, "Application of coding theory to the construction of modular lattices", *J. Combin. Theory Ser. A*, Vol. 78, pp. 92-119, 1997.
- [2] K. Betsumiya, S. Georgiou, T.A. Gulliver, M. Harada and C. Koukouvinos, "On self-dual codes over some prime fields", *Discrete Math*, vol 262, pp. 37–58, 2003.
- [3] W. Bosma, J. Cannon and C. Playoust, "The Magma algebra system. I. The user language", *J. Symbolic Comput.*, vol. 24, pp. 235—265, 1997.
- [4] J. H. Conway, N. J. A. Sloane, "A new upper bound on the minimal distance of self-dual codes", *IEEE Trans. Inform. Theory*, Vol. 36, 6, 1319–1333, 1990.
- [5] S.T. Dougherty, P. Gaborit, M. Harada and P. Solé, "Type II codes over $\mathbb{F}_2 + u\mathbb{F}_2$", *IEEE Trans. Inform. Theory*, Vol. 45, pp.32–45, 1999.
- [6] S.T. Dougherty, T.A. Gulliver, M. Harada, "Extremal binary self dual codes", *IEEE Trans. Infrom. Theory*, Vol. 43, No.6, pp. 2036–2047, 1997.
- [7] S.T. Dougherty, J.L. Kim, H. Kulosman and H. Liu, "Self-dual codes over commutative Frobenius rings", *Finite Fields Appl.*, Vol.16, pp.14–26, 2010.
- [8] S.T. Dougherty, B. Yildiz and S. Karadeniz, "Codes over R_k, Gray Maps and their Binary Images", *Finite Fields Appl.* Vol. 17, No. 3, pp.205–219, 2011.
- [9] S.T. Dougherty, B.Yildiz and S.Karadeniz, "Self-Dual Codes over R_k and Binary Self-Dual Codes", *Eur. J. Pure Appl. Math.*, Vol.6, No.1, pp. 89-106, 2013.
- [10] S. Karadeniz, B. Yildiz, "New extremal binary self-dual codes of length 64 from R_3 lifts of the extended binary Hamming code", *Des. Codes Cryptogr.,* Vol. 74, No. 3, pp. 673–680, 2015.
- [11] S. Karadeniz, B. Yıldız and N. Aydın, "Extremal binary self-dual codes of lengths 64 and 66 from four-circulant constructions over codes $\mathbb{F}_2 + u\mathbb{F}_2$", *FILOMAT*, Vol. 28, Issue 5, pp. 937-945, 2014.
- [12] A. Kaya, B. Yildiz, İ. Şiap, "New extremal binary self-dual codes from $\mathbb{F}_4 + u\mathbb{F}_4$-lifts of quadratic double circulant codes over $\mathbb{F}_4$", *Finite Fields Appl.*, Vol.35, pp.318–329, 2015.
- [13] A. Kaya, B. Yildiz, "Various constructions for self-dual codes over rings and new binary self-dual codes", to appear in *Discrete Mathematics.*
- [14] A. Kaya, B. Yıldız, "New extremal binary self-dual codes of length 68", *Journal of Algebra Combinatorics Discrete Structures and Applications*, Vol. 1, No. 1, pp 29-39, 2014.
- [15] A. Kaya, N. Tufekci, "New extremal binary self-dual codes of lengths 66 and 68 from codes over $R_{k,m}$", available online at http://arxiv.org/abs/1502.06945
- [16] A. Kaya, B. Yıldız and A. Pasa, *Binary generator matrices of new extremal self-dual binary codes of lengths 64, 66 and 68*, available online at http://www.fatih.edu.tr/~aserdogan/kyp.html
- [17] E. M. Rains, "Shadow Bounds for Self Dual Codes", *IEEE Trans. Inf. Theory*, Vol.44, pp.134–139, 1998.
- [18] N. Yankov, "Self-dual $[62, 31, 12]$ and $[64, 32, 12]$ codes with an automorphism of order 7", *Advances in Mathematics of Communications*, Vol.8, No1. pp.73–81, 2014.
- [19] N. Yankov, M.-H. Lee, M. Gurel, M. Ivanova, "Self-dual codes with an automorphism of order 11", *IEEE Trans. Inform. Theory*, Vol. 61, No. 3, pp. 1188–1193, 2015.
- [20] B. Yildiz and S. Karadeniz, "Linear codes over $\mathbb{F}_2+u\mathbb{F}_2+v\mathbb{F}_2+uv\mathbb{F}_2$", *Des. Codes Cryptogr.*, Vol. 54, No. 1, pp.61–81, 2010.

Department of Mathematics, Fatih University, 34500 Istanbul, Turkey
E-mail address: `nabidin@gmail.com,byildiz@fatih.edu.tr,abdullah.pasa@fatih.edu.tr`